


 Ecole d'ingénieurs


 INA P-G


 INRA

Fate of isoxaflutole and its diketonitrile metabolite
under conventional and conservation tillage in an
irrigated continuous-maize field

L. Alletto^{1,2}, Y. Coquet² & C. Labat²




¹ École d'ingénieurs de Purpan, Agronomy department, Toulouse, France.

² UMR INRA/INA PG Environment and Arable Crops, Thiverval-Grignon, France.


 Agence de l'Eau
Adour Garonne

Diffuse Inputs into the Groundwater – January 29-31th, 2007 - Graz

Outline



Introduction

- Context of the study
- The cropping systems
- Objectives of the study

Materials and methods

- The experimental site: localisation and soil characteristics
- Isoxaflutole (IFT) properties
- Sampling procedure: soil
- Sampling procedure: water

Results and discussion

- General data
- Persistence of isoxaflutole
- Water and herbicide leaching

Conclusion

2
PURPAN

Introduction


Context

National context: generalization of water resources pollution by pesticides...

Economic context: Midi-Pyrénées region = 2nd region for maize production in France

Agronomic context: Typical maize production management in the region:

- 80 % of the production is in continuous maize with more than 60 % irrigated.
- Tillage usually included a mouldboard ploughing (30-cm depth) at the end of the winter
- Soils are unprotected during the inter-crop (from November to May)



Environmental context:


⇒ In the region, this system of production has generated several environmental problems (nitrate, atrazine)


⇒ Now atrazine is forbidden: there is a need for development of new strategies to control weeds in continuous maize systems

3
PURPAN

Introduction


Context





Conventional tillage (CT)

/




Conservation tillage (MT)

vs. Pesticide ?

- Organic carbon content ↑ surface → pesticides sorption ↑ (Locke et al., 1997)
Desorption: very few studies... tend to increase under MT (Ding et al., 2002)
- Degradation: highly contrasted results !
- Runoff and erosion: MT are efficient to reduce erosion, but runoff depends on climatic conditions (Fawcett et al. 1994)
- Leaching: contrasted results but for no-tillage systems leaching of pesticides increases (Watts & Hall, 1996)

There is a need to evaluate and/or design new cropping systems to both maintain weed control efficiency and limit environmental impacts


4
PURPAN



Introduction

The cropping systems: subsurface differences


Conventional tillage (CT)



Labels in CT image: A clod, Plough pan, Crop residues, F clod, Inter-furrow.

(Manichon, 1982; Roger-Estrade et al., 2004)

Conservation tillage (MT)



Labels in MT image: No visible limit of the previously ploughed layer, Root development > 70 cm, Undulations created by the disc harrowing, Vertical cracks.

→ Consequences on water dynamics and solutes transport


7 PURPAN

Introduction



Objectives of the study

Evaluate the effects of tillage practices (conventional vs. conservation tillage ; inter-crop with or without cover-crop) on:

- 1- Isoxaflutole degradation and formation of diketonitrile
- 2- Leaching potential of isoxaflutole and its diketonitrile metabolite




8



Isoxaflutole (IFT) properties

- Proherbicide, isoxazoles



- Annual grasses and broadleaves weeds, roots uptake

- Pre-emergence of maize (75 g ha⁻¹)

- Inhibitor of the biosynthesis of carotenoids

- IFT: low solubility** in water (6.2 mg L⁻¹), **rapidly degraded** (DT_{50} : 1.4-3 j), **good retention** on organic compounds (K_{OC} : 122 L kg⁻¹)

- Degradation: formation of the **diketonitrile (DKN)** with a **higher solubility** (300 mg L⁻¹), a **lower retention** (K_{OC} : 92 L kg⁻¹) and a **higher persistence** (DT_{50} : 8-16 j)




11

PURPAN

Sampling procedure: soil

- Before treatment: sampling of soil from surface to 80-cm depth to control initial concentration of IFT and DKN
- Treatment day: control of the variability of the treatment with fibreglass paper
- Sampling of soil from surface to a maximum of 30-cm depth
- Sampling time: t_{ini} , t_0 , t_2 , t_3 , t_5 , t_7 , t_{11} , t_{14} , t_{21} , t_{28}
- Storage of frozen samples (-18°C) until analysis
- Analysis by HPLC-MS/MS



12

PURPAN

Sampling procedure: water

- Ceramic cups: 4 by soil pit at 20 and 70 cm-depth
- Fibreglass wick lysimeters: 2 by soil pit at 40 cm (25x25 cm). Fibreglass wick length: 70 cm
- Sampling of soil water with ceramic cups and fibreglass wick lysimeters
- Storage of frozen samples (-18°C) until analysis
- Analysis by HPLC-MS/MS

Materials-methods

13

PURPAN

General data

- In soil samples: Limit of quantification (LOQ) $\approx 0.01 \text{ mg a.i. kg}^{-1} \text{ soil}$
- In water samples: LOQ depends on collected volumes (V)
 - If $V < 50 \text{ ml} \rightarrow \text{LOQ} = 0.2 \mu\text{g L}^{-1}$
 - If $V > 1000 \text{ ml} \rightarrow \text{LOQ} = 0.02 \mu\text{g L}^{-1}$

In 2005:

- **400 soil samples** were analysed from treatment day to 28 DAT
- **73 water samples**: (14 from the ceramic cups and 59 from the fibreglass wick lysimeters)



Rainfall data during the cropping season

- Total precipitation: **599 mm**
- First rainfall: **1 DAT – 7.6 mm in 6 h**

Results-discussion

Total : 599 mm
 - rainfall : 255 mm
 - irrigation : 343 mm

14



- Tillage practices had no effect on the in-field degradation of isoxaflutole
- Residues on soil surface seemed to slow down degradation rate of IFT

Effect of interception and retention ?

- Migration in soil was faster and more important under conventional technique (data not shown)
- Water drainage was two times higher under conventional technique
- Herbicide leaching was between 2 and 7 times lower under conservation technique, with the lowest leaching under the cover-crop plot

Retention processes and/or degradation were modified by tillage and residues management ?

Thank you.

